

A Proposed Culinary Medicine Education to Strengthen Medical Nutrition Therapy for Improve Health Outcomes in Cirrhosis



Daniella Hernandez BS, Ashley Mullins MS, RD, LD, CNSC

Department of Nutritional Sciences | Texas Christian University

BACKGROUND

Cirrhosis is the advanced stage of chronic liver injury marked by progressive fibrosis and hepatic dysfunction resulting from alcohol-associated liver disease, chronic viral hepatitis, or metabolic dysfunction. It may ultimately progress to decompensation with complications such as portal hypertension, ascites, variceal bleeding, and hepatic encephalopathy.¹⁻² Malnutrition and sarcopenia are highly prevalent (up to 70%) in cirrhosis and are associated with increased hospitalization, infection risk, and mortality.³ Medical nutrition therapy (MNT) is a cornerstone of cirrhosis management that includes alcohol cessation, adequate energy intake, increased protein intake, sodium restriction, and avoidance of prolonged fasting.^{4,5} Despite clear guidelines, adherence is often limited by practical ability to translate clinical recommendations into sustainable daily eating patterns. This case report offers culinary medicine (CM) as a preventative, skill-based intervention to reduce progression from compensated to decompensated cirrhosis by improving nutrition-related behavior change and dietary adherence. Based on established guidelines, three culinary medicine targets are proposed: (1) supporting alcohol cessation through structured meal timing, (2) implementing sodium-reduction strategies utilizing herbs, and spices to preserve palatability, and (3) optimizing protein distribution across meals and snacks to mitigate sarcopenia and reduce catabolic stress. Integrating clinical nutrition guidelines with culinary skills may empower patients and reduce risk of cirrhosis-related complications.

NUTRITIONAL CONSIDERATIONS

Translating clinical guidelines into practical food-based strategies remains essential for supporting implementation in clinical and community settings. **Table 1** summarizes evidence-based MNT recommendations alongside related culinary medicine applications for patients who have cirrhosis.

Table 1: Cirrhosis MNT Recommendations with Culinary Medicine Application

Nutritional Considerations	Evidence-Based MNT Recommendation ^{3, 6-8}	Culinary Medicine Application ⁹⁻¹⁶
Energy and Protein Requirements	- Increased metabolic demand - Energy: 30-35 kcal/kg/day - Protein: 1.2-1.5 g/kg/day - Avoid protein restriction unless in severe encephalopathy	- Incorporate protein at each meal and snack to support daily intake distribution - Build balanced meals using affordable and sustainable protein sources
Meal Timing and Fasting Avoidance	- Reduced glycogen stores increase early fasting catabolism - Frequent meals every 3-4 hours - Late-evening snacks (~200 kcals) improves nitrogen balance and limits muscle loss	- Establish structured meal routines throughout the day - Plan evening snacks - Provide simple snack preparation strategies
Sodium Management	Sodium restriction ≤ 2,000 mg/d for ascites management	- Flavor enhancement using sodium-free herbs and spices - Culinary techniques emphasizing aromatics and acids
Alcohol Cessation and Nutrition Recovery	- Alcohol cessation is essential for disease stabilization - Refer for additional support	- Structured meal timing using behavioral anchors replacing drinking routines - Develop alternative beverage habits

CASE REPORT

Summary:

- 64 YO F with alcohol-associated cirrhosis (diagnosed ~11 years) admitted for acute chest pain and found to have severe aortic stenosis requiring an aortic valve replacement (AVR)
- Developed abdominal distention; evaluation confirmed ischemic colitis with non-functional gastrointestinal (GI) tract
 - Underwent surgical intervention resulting in bowel discontinuity and inability to initiate enteral nutrition (EN)
 - Hospitalization complicated by GI bleeding, septic shock, respiratory failure requiring intubation, and acute kidney injury requiring continuous renal replacement therapy (CRRT)
 - Progressed to multi-organ failure and died on hospital day 33

At this stage, the patient was unable to participate in nutrition-related education or behavior change due to critical illness, altered cognition, and lack of oral intake.

TIMING OF CULINARY MEDICINE INTERVENTION

Compensated Cirrhosis

✓ Highest opportunity for CM intervention

Focus:

- Establish sustainable nutrition behaviors
- Translate MNT into practical daily routines
- Support long-term disease management
- Refer for additional support & counseling

Patients are physically and cognitively able to engage in behavior change and skill development

Early Decompensation

✓ CM intervention remains feasible

Focus:

- Maintain energy and protein intake
- Simplify meal preparation to reduce burden of change
- Reinforce structured meal & snack timing

Barriers begin to arise that include fatigue, early satiety, low motivation, decreased appetite, and increased symptoms

Decompensated Cirrhosis

X Not appropriate for CM intervention

Focus:

- Life-sustaining medical and nutrition support

Why CM is Not Appropriate:

- Unable to participate in education
- Cognitive impairment
- Acute instability
- Limited or absent oral intake (NPO, EN, or TPN)

At this point, nutrition care shifts to critical care support

REFERENCES

1.Cleveland Clinic. Cirrhosis of the Liver. Cleveland Clinic. July 26, 2023. Accessed November 1, 2025. <https://my.clevelandclinic.org/health/diseases/15572-cirrhosis-of-the-liver>

2.Sharma B, John S. Hepatic Cirrhosis. National Library of Medicine. October 31, 2022. Accessed November 1, 2025. <https://www.ncbi.nlm.nih.gov/books/NBK482419/>

3.Lai JC, Tandon P, Bernal W, et al. Malnutrition, frailty, and sarcopenia in patients with cirrhosis: 2021 practice guidance by the American Association for the Study of Liver Diseases. *Hepatology*. 2021;74(3):1611-1644. doi:10.1002/hep.32049

4.European Association for the Study of the Liver. EASL clinical practice guidelines on nutrition in chronic liver disease. *J Hepatol*. 2019;70(1):172-193. doi:10.1016/j.jhep.2018.06.024

5.Moss O. Nutrition Priorities: Diet recommendations in liver cirrhosis. *Clin Liver Dis (Hoboken)*. 2019;14(4):146-148. doi:10.1002/cld.831

6.Plauth M, Bernal W, Dasarthy S, et al. ESPEN guideline on clinical nutrition in liver disease. *Clinical Nutrition (Edinburgh, Scotland)*. 2019;38(2):485-521. doi:10.1016/j.clnu.2018.12.022

7.Fishman G, Singer P. Metabolic and nutritional aspects in continuous renal replacement therapy. *Journal of Intensive Medicine*. 2023;3(3):228-238. doi:10.1016/j.jointm.2022.11.001

8.da Silva JSV, Seres DS, Sabino K, et al. ASPEN consensus recommendations for refeeding syndrome. *Nutrition in Clinical Practice*. 2020;35(2):178-195. doi:10.1002/ncp.1047

9.Mamerow MM, Mettler JA, English KL, et al. Dietary protein distribution positively influences 24-h muscle protein synthesis in healthy adults. *J Nutr*. 2014;144(6):876-880. doi:10.3945/jn.113.185280

10.Chen C, Wang L, Kuo H, Fang Y, Lee H. Significant effects of late evening snack on liver functions in patients with liver cirrhosis: A meta-analysis of randomized controlled trials. *J Gastroenterol Hepatol*. 2019;34(7):1143-1152. doi:10.1111/jgh.14665

11.Res PT, Groen B, Pennings B, et al. Protein ingestion before sleep improves postexercise overnight recovery. *Med Sci Sports Exerc*. 2012;44(8):1560-1569. doi:10.1249/MSS.0b013e31824cc363

12.Anderson CA, Cobb LK, Miller ER, et al. Effects of a behavioral intervention that emphasizes spices and herbs on adherence to recommended sodium intake: Results of the SPICE randomized clinical trial. *Am J Clin Nutr*. 2015;102(3):671-679. doi:10.3945/ajcn.114.100750

13.Douglas A, Vannereux M, Giboreau A. The impact of herbs and spices on increasing the appreciation and intake of low-salt legume-based meals. *Nutrients*. 2019;11(12):2901. doi:10.3390/nu11122901

14.Ting PS, Lin WT, Liangpunsakul S, et al. Convergence of alcohol consumption and dietary quality in US adults who currently drink alcohol: An analysis of two core risk factors of liver disease. *Nutrients*. 2024;16(22):3866. Published 2024 Nov 13. doi:10.3390/nu16223866

15.Butler L, Popkin BM, Poti JM. Associations of alcoholic beverage consumption with dietary intake, waist circumference, and body mass index in US adults: National Health and Nutrition Examination Survey 2003-2012. *J Acad Nutr Diet*. 2018;118(3):409-420.e3. doi:10.1016/j.jand.2017.09.030

16.Breslow RA, Guenther PM, Juan W, Graubard BI. Alcoholic beverage consumption, nutrient intakes, and diet quality in the US adult population, 1999-2006. *J Am Diet Assoc*. 2010;110(4):551-562. doi:10.1016/j.jada.2009.12.026

CULINARY MEDICINE STRATEGIES WITH CIRRHOSIS

Target 1: Protein Distribution and Snack Timing to Reduce Catabolic Stress⁹⁻¹¹


- Even distribution of protein intake across meals supports greater daily muscle protein synthesis
- Minimizing prolonged fasting through planned snacks, particularly late-evening intake to reduce overnight catabolic stress observed in cirrhosis populations
- Structured protein timing represents a feasible nutrition strategy to sustain overnight amino acid availability and stimulate protein synthesis and mitigate sarcopenia risk in chronic liver disease

Target 2: Sodium Restriction While Maintaining Palatability^{12,13}

- Flavor-enhancement strategies using sodium-free herbs and spices to maintain food acceptability and overall meal satisfaction leading to long-term dietary adherence
- Providing CM education classes that translate sodium restriction guidelines into practical cooking behaviors by teaching flavor substitution strategies rather than simply advising patients to reduce salt intake

Target 3: Structured Meal Timing as a Behavioral Anchor to Reduce Alcohol Consumption¹⁴⁻¹⁶

- Alcohol consumption is consistently associated with poor dietary quality and altered eating behaviors, identifying nutrition patterns as intervention points
- Establishing structured meal timing and replacing alcohol with nutrient-dense alternative beverages may provide stability that supports nutritional adequacy
- Shared meals with family or peers provide social support and promote more consistent eating patterns



CONCLUSION

Cirrhosis management relies heavily on nutrition strategies that patients must sustain outside of the clinical setting, yet translating MNT into daily eating behaviors remains challenging. This case demonstrates that culinary medicine provides a feasible, skill-based framework for translating structured meal timing, sodium reduction techniques, and strategic protein distribution into sustainable eating practices. The patient’s clinical course highlights the challenges patients’ face when nutrition recommendations are difficult to translate. Culinary medicine represents a scalable and practical approach for dietitians and healthcare professionals to operationalize nutrition guidelines through everyday food preparation, helping patients consistently apply practical nutrition recommendations in their daily lives.